DRIFTING IN THE ABYSS

A Comprehensive Interdisciplinary Analysis of Rogue Planets with Hidden Subsurface Oceans, Geothermal Biospheres, and Cosmic Demographics

PREPARED BY	Gemini Interdisciplinary Research Engine Planetary Science & Astrobiology Division
DATE OF MODELS	June 13, 2026
DOCUMENT ID	RP-H2O-DEEP-2026-V1
CLASSIFICATION	Unrestricted Scientific Synthesis

Executive Summary & Core Hypothesis

The classical definition of planetary habitability has historically been inextricably tethered to the concept of a host star's Circumstellar Habitable Zone (CHZ)—the spatial band where solar irradiance maintains liquid water on a terrestrial world's surface. However, the discovery and theoretical framework of free-floating planetary-mass objects, or **rogue planets**, challenges this stellar-centric paradigm. Ejected from their birth systems via chaotic multi-body dynamical interactions or formed through direct isolation-mode gravitational collapse, these worlds drift through the near-absolute zero thermal environment of interstellar space.

> **THE STARLESS THERMODYNAMIC EQUATION**
>
> ***No Sun + Internal Energy + Subsurface Water → Potential Habitable Environment***

This analysis delivers a comprehensive investigation into the mechanics of starless planetary architectures. It explores how internal radiogenic decay, tidal friction, and primordial primordial thermodynamic reserves can provide a robust, long-term heat budget. When shielded by a dense, collision-induced absorption (CIA) hydrogen atmosphere or an insulating cryogenic lithospheric ice shell, this thermal flux is sufficient to preserve extensive liquid water oceans beneath hundreds of kilometers of solid ice. This interdisciplinary paper synthesizes astrophysics, planetary geology, extremophile microbiology, and information theory to map out the thermodynamic boundaries, biochemistry, evolutionary trajectories, and technological capacities of life inside these dark, ubiquitous worlds.

1. Astrophysical Origins & Galactic Demographics

Rogue planets inhabit the interstellar medium, decoupled from the gravitational governance of any stellar host. Planetary demographic surveys reveal that these objects are not rare anomalies, but rather primary products of the stellar and planetary formation processes.

1.1 Dynamic Ejection Pathways

The most prevalent mechanism generating low-mass rogue planets is dynamical scattering during the early, chaotic phases of solar system formation. In young, gas-rich protoplanetary discs, runaway core accretion produces multiple giant planets alongside smaller terrestrial protoplanets. As these bodies migrate, they enter orbital resonance chains. If the architecture becomes unstable, giant planets undergo close gravitational encounters, inducing sudden, severe changes in eccentricity and semi-major axes.

During these events, light terrestrial planets or low-mass gas giants are dynamically accelerated beyond the system's escape velocity. Recent physical simulations demonstrate that binary star systems, which account for roughly 40% of all stellar systems, possess an exceptionally high cross-section for planetary ejection via a violent "pincer movement" generated by the complex, rotating gravitational fields of the two host stars (Balbi et al., 2020).

1.2 Isolated Starless Formation (Direct Collapse)

An alternative evolutionary vector involves direct gravitational collapse within giant molecular clouds. This mechanism operates identically to standard star formation but occurs at the extreme sub-stellar end of the IMF (Initial Mass Function). If a dense clump of interstellar gas experiences turbulent fragmentation, it can surpass its local Jeans mass without possessing sufficient material to form a brown dwarf or low-mass red dwarf. The resulting objects cross the sub-brown dwarf boundary (typically defined around $13\ M_{Jup}$, the threshold for deuterium fusion) and form starless planetary-mass objects directly within star-forming clusters.

1.3 Galactic Abundances and Population Architecture

Quantifying the precise space-density of rogue worlds is an active focus of modern wide-field gravitational surveys. Theoretical population synthesis models indicate that the interstellar medium contains a vast, hidden demographic of nomadic objects (Strigari et al., 2012). Ground-based and space-based microlensing events establish that low-mass rogue planets (ranging from Mars to Earth mass) are significantly more common than giant gas-giant nomads.

Statistical extrapolations assert that rogue planets likely outnumber stars in the Milky Way by orders of magnitude, implying that the average distance between a random observer in interstellar space and the nearest rogue world is less than a few light-years.

Classification	Mass Range	Primary Origin Mechanism	Atmospheric/Surface Profile
Sub-Brown Dwarfs	1 - 13 M_{Jup}	Molecular Cloud Fragmentation	Thick H/He envelopes, high primordial temperatures
Jovian Nomads	0.5 - 5 M_{Jup}	Gas-Giant Resonance Scattering	Dense volatile layers, intense residual heat
Super-Earth / Terrestrial	0.1 - 10 $M_{\oplus}$	Planet-Planet Gravitational Scattering	Cryo-atmospheres, volatile ice shells, rocky cores

2. Internal Energy Sources & Geothermal Budgets

Deprived of stellar insolative flux, a rogue planet's surface equilibrium temperature drops to match the cosmic microwave background (~3 K) modified by galactic cosmic rays and starlight, resulting in an effective background temperature of ~10–40 K. To maintain liquid water, the planet must rely entirely on its interior thermodynamic engine.

2.1 Long-Lived Radiogenic Decay

The foundational core energy source for any terrestrial-type rogue world is the radioactive decay of unstable, long-lived lithophilic isotopes within its mantle and crust. The primary elements driving this process are Uranium-238 (^{238}U), Uranium-235 (^{235}U), Thorium-232 (^{232}Th), and Potassium-40 (^{40}K).

On Earth, this radiogenic mechanism generates approximately 20 Terawatts (TW) of the total 46 TW surface heat flux. For a rogue planet, the total radiogenic heat generation over time can be tracked using the integrated decay law:

$$Q(t) = \Sigma\, C_i \bullet H_i \bullet e^{-\lambda_i t}$$

Where C_i represents the concentration of isotope i at formation, H_i is the specific heat release per unit mass, and λ_i is the decay constant. Super-Earth class rogue worlds (3 to 10 Earth masses) with iron-silicate structures carry substantially larger radiogenic inventories. Due to their lower surface-area-to-mass ratio, they exhibit significantly slower thermal dissipation rates, extending the longevity of their geodynamo and core thermal output over tens of billions of years.

2.2 Exo-Moon Tidal Dissipation

While it may seem intuitive that an ejected planet drifts alone, hydrodynamical scattering simulations demonstrate that gas giants and massive Super-Earths frequently retain their tightly bound satellite architectures during ejection events. A rogue planet flanked by an orbital system of moons experiences continuous gravitational distortions.

If the orbits are forced to maintain non-zero eccentricities due to mutual satellite resonances (analogous to the Laplace resonance between Io, Europa, and Ganymede), the continuous cyclic distortion of the planet's body generates enormous internal frictional shear. This tidal heating dissipation can be modeled by:

$$E_{tidal} = -\,(21 / 2) \bullet (k_2 / Q) \bullet (G\, M_{sm}^{\,2}\, R^5\, n\, e^2) / a^6$$

Where k_2 is the planet's potential Love number, Q is the tidal dissipation quality factor, M_{sm} is the mass of the satellite, a is the semi-major axis, n is the mean orbital motion, and e is the eccentricity. Frictional shear transforms this kinetic orbital energy directly into thermal flux at the core-mantle interface, creating localized volcanic zones and active hydrothermal systems.

2.3 Residual Formation & Differentiation Energy

During accretion, gravitational potential energy is converted directly into heat, melting the young planet's interior and inducing core differentiation (the sinking of heavy iron-nickel phases to the interior). This primordial heat is trapped deep within the mantle.

The cooling timescale of a solid planet scales quadratically with its radius:

$$\tau \propto R^2 / \kappa$$

Where κ is the thermal diffusivity of the mantle rock. For massive Super-Earths, this ensures that even in the total absence of external stellar radiation, the primordial heat of formation can take 5 to 10 billion years to escape, providing a stable baseline flux.

3. The Hidden Ocean Hypothesis & Stratigraphy

The concept of a subglacial ocean on a starless world requires an intricate thermodynamic balance where the upward geothermal heat flux balances the conductive heat loss through a thick crystalline outer boundary layer. This class of worlds is often categorized as "Steppenwolf planets" (Abbot & Switzer, 2011).

3.1 Macro-Structural Layering

The internal architecture of a habitable rogue ocean world typically consists of five highly distinct stratigraphical zones:

1. Cryo-Atmosphere / Surface Glacial Rind: An outer shell composed of hard, vacuum-temperature water ice, combined with condensed volatile frosts (nitrogen, carbon dioxide, methane) at temperatures under 40 K.

2. Conductive Ice I Shell: A thick layer of standard crystalline hexagonal ice. Heat transfer through this layer is strictly conductive, maintaining a sharp linear thermal gradient from ~40 K at the surface to 273 K at the interior base.

3. Subsurface Liquid Ocean: A global aqueous layer sustained by internal heat. The depth of this ocean can span from tens to hundreds of kilometers, dwarfing Earth's hydrosphere.

4. High-Pressure Polymorph Ice Floor: On worlds with masses exceeding 2–3 Earth masses or containing very high water-mass fractions, the hydrostatic pressure at the base of the ocean exceeds several gigapascals. This forces liquid water to freeze into exotic high-density solid phases, such as Ice V, Ice VI, or Ice VII.

5. Silicate Mantle & Core: The rocky basement of the planet, rich in radiogenic elements and driving volcanism or serpentinization reactions.

> *"The critical structural constraint for life is whether a high-pressure ice floor physically uncouples the liquid ocean from the underlying rocky silicate mantle. On smaller worlds or those with low water budgets, the ocean sits directly on a rocky crust, enabling rich water-rock chemical reactions. On hyper-water-rich Super-Earths, the ocean is sandwiched between two ice layers, constraining chemical transport to fracture networks and convective plumes."*

3.2 Hydrothermal and Geochemical Interfaces

Where liquid water makes contact with the silicate crust, high-temperature fluid circulation drives extensive alteration chemistry. Seawater filtering through basaltic fissures reacts with olivine-rich minerals in a process known as **serpentinization**:

$$3\,Fe_2SiO_4 + 2\,H_2O \rightarrow 2\,Fe_3O_4 + 3\,SiO_2 + 2\,H_2$$

This exothermic reaction generates large quantities of magnetite, silica, and molecular hydrogen (H_2). The release of hydrogen, paired with primordial dissolved carbon dioxide, establishes a highly potent geochemical non-equilibrium state, generating a steady flux of chemical energy that can substitute for solar photon flux.

4. Atmospheres Without a Star: Thermal Insulation

A rogue planet's capability to retain an unfrozen internal ocean is immensely elevated if it can maintain a thick gas envelope capable of delaying core thermal leakage.

4.1 Primordial Hydrogen Greenhouse Engines

During the accretion phase inside a gas-rich circumstellar disc, rocky protoplanets quickly accumulate dense atmospheres derived from the surrounding solar nebula, dominated by molecular hydrogen (H_2) and helium (He). For planets bound to a bright host star, this primordial hydrogen envelope is rapidly stripped away via extreme ultraviolet (EUV) photoevaporation and stellar wind stripping (Zhang, 2020).

Rogue planets, however, escape this stellar erosion vector. If a world is scattered out of its system early, its massive hydrogen envelope remains intact. While symmetric diatomic molecules like H_2 do not normally act as greenhouse gases on Earth, the extreme pressures characteristic of a primordial envelope (exceeding 100 to 1000 bars) induce a phenomenon known as **Collision-Induced Absorption (CIA)**.

When gas density is exceptionally high, frequent intermolecular collisions briefly distort the electron clouds of colliding H_2-H_2 or H_2-He pairs, creating transient dipole moments. These dipoles absorb heavily across the far-infrared spectral bands, precisely where the planet's internal geothermal heat attempts to radiate into space. This mechanism acts as an ultra-efficient thermal blanket.

4.2 Atmospheric Preservation and Volatile Collapse

For lighter worlds lacking the gravity to hold light hydrogen, heavier volatile gases like carbon dioxide (CO_2), nitrogen (N_2), and methane (CH_4) are released via ongoing volcanic outgassing. However, in the absence of a star, these gases face the threat of *atmospheric collapse*.

As the surface cools below the condensation point of these compounds, they precipitate out as solid snow, forming massive global glaciers. Yet, physical models demonstrate that a balance can be achieved (Abbot & Switzer, 2011). As the gas condenses into a surface ice sheet, the ice sheet itself acts as a powerful solid-state thermal insulator. The geothermal heat flux passing up from the core warms the lower boundary of the ice sheet, recreating an active, high-pressure subglacial gas envelope or melting the ice directly from below to sustain the ocean layer.

5. Energetics & Energetic Chemistry of Starless Biospheres

Because rogue oceans exist under total optical darkness beneath hundreds of kilometers of ice and gas, photosynthesis via chlorophyll is physically impossible. Any biological architecture must operate using chemosynthesis.

5.1 Chemotrophic Metabolic Pathways

The base of a starless biosphere must rely entirely on primary producers that exploit inorganic chemical energy gradients. The primary metabolic pathways available within these dark liquid oceans are summarized below:

1. Autotrophic Methanogenesis: Utilizing the molecular hydrogen generated by serpentinization alongside dissolved carbon dioxide:

$$CO_2 + 4\,H_2 \rightarrow CH_4 + 2\,H_2O \quad [\Delta G^\circ = -131\ kJ/mol]$$

2. Sulfate Reduction: Utilizing sulfate minerals dissolved from hydrothermal leaching to oxidize volcanic gases:

$$SO_4^{2-} + 4\,H_2 + 2\,H^+ \rightarrow H_2S + 4\,H_2O$$

3. Iron Oxidation: Microbes utilizing dissolved iron species (Fe^{2+}) exported via hydrothermal vent fields as electron donors to fix inorganic carbon into cellular mass.

5.2 Analogous Earth Systems and Energy Limits

We possess strong empirical proof for the validity of these dark biospheres from analog environments on Earth. The deep lithospheric biosphere, extending several kilometers into Earth's crust, hosts massive microbial populations that are completely decoupled from surface sunlight (Schulze-Makuch & Fairén, 2021).

Similarly, ecosystems surrounding submarine hydrothermal vents (such as the alkaline vent fields of the Lost City) demonstrate that complex multi-tiered biological webs can thrive exclusively on chemical energy. However, because the total energy flux from internal geothermal sources is roughly three to four orders of magnitude smaller than the solar irradiance striking Earth's surface, the absolute biological biomass of a rogue planet will be severely energy-limited, maintaining a low-density, highly efficient ecological architecture.

6. Origin of Life: Abiogenesis vs. Ejection Survival

A critical question in astrobiology is whether life can independently originate on a starless world, or if it must be inherited from a circumstellar past.

6.1 Scenario A: Pre-Ejection Circumstellar Development

In this scenario, abiogenesis occurs early within the planet's native solar system, while it is still bound to its parent star. Life populates the surface oceans via standard solar-driven primordial mechanisms. During the gravitational scattering and subsequent ejection phase, the surface ocean completely freezes over into an insulating ice shell over a timescale of 10^4 to 10^6 years.

While surface life is completely eradicated, deep microbial populations situated near hydrothermal vents or inside deep porous basalt aquifers experience virtually no disruption. The ice shell acts as a shield against the intense cosmic radiation of interstellar space, preserving a pre-existing biosphere (Sadlok, 2020).

6.2 Scenario B: In-Situ Dark Abiogenesis

Alternatively, life can emerge directly on a world drifting through interstellar space. Recent laboratory work in prebiotic chemistry shows that the fundamental building blocks of life do not require UV light to synthesize. Deep, porous alkaline hydrothermal vents act as natural electrochemical flow reactors.

The walls of these microscale vents, composed of iron-sulfide minerals like mackinawite (***FeS***), act as powerful inorganic catalysts. They possess structures strikingly similar to the active sites of modern metabolic enzymes (such as ferredoxins). The steep pH gradient between the alkaline hydrothermal fluid (pH ~9–11) and the acidic, CO_2-rich ocean water (pH ~5–6) creates a continuous proton-motive force across these mineral walls, driving the synthesis of amino acids and RNA polymers in complete darkness.

7. Detectability & Observational Methodologies

Because rogue ocean planets emit no reflected light and are located light-years away from any stellar illumination source, discovering them presents an extraordinary challenge for observational astronomy.

7.1 Space-Based Gravitational Microlensing

The most effective technique to confirm the existence and mass function of rogue planets is gravitational microlensing. When a rogue planet passes directly along the line of sight between a distant background star and a space telescope, its gravitational field acts as a natural lens, bending and magnifying the background star's light.

The duration of this magnification spike is directly proportional to the mass of the lens object. While a stellar lens produces a signal lasting weeks, a terrestrial rogue planet creates a brief spike lasting only a few hours to less than two days. Modern space observatories, such as the Nancy Grace Roman Space Telescope's Galactic Bulge Time-Domain Survey and the Euclid telescope, use high-cadence near-infrared sensors to monitor millions of stars simultaneously, logging hundreds of these starless planetary events.

7.2 Far-Infrared and Microwave Thermal Signatures

To verify if a rogue planet hosts a hidden ocean, astronomers must observe its direct thermal emissions. Because a habitable rogue world is warmed internally, it acts as a faint blackbody radiator emitting in the far-infrared and microwave bands (***50 - 200 μm***).

By measuring the precise spectral energy distribution (SED), researchers can map out the planet's effective temperature. If the world is a dead, solid block of rock and ice, its surface heat distribution will match simple conductive transport models. However, if it contains a convective subsurface liquid ocean or active tidal plume venting, the thermal emission will reveal distinct anomalies and trace volatile signatures (such as water vapor, methane, or ammonia plumes) detectable during exceptionally close interstellar flybys or via sub-millimeter array spectroscopy.

8. Evolutionary Consequences of Dark Biospheres

If life successfully persists over billions of years within an isolated, starless ocean, it will experience evolutionary pressures fundamentally distinct from anything observed on Earth.

8.1 Complete Absence of Phototropism

On Earth, the biological clock, sensory apparati, and behavioral cycles of nearly all organisms are coordinated by the diurnal solar cycle. In a starless biosphere, the concept of vision via visible light is completely absent. Sensory evolution will instead favor extreme specialization in alternative physics:

Mechanoreception & Acoustic Navigation: Because water under high pressure is an exceptional acoustic conductor, organisms will utilize active and passive echolocation to map their three-dimensional aquatic environments, adapting to sound waves reflecting off the upper ice ceiling and lower rock floor.

Chemotaxis & Olfactory Arrays: Complex chemical sensing networks will allow organisms to track concentration gradients of vital nutrients, tracking hydrothermal plumes across thousands of kilometers.

Electrosensation: Much like Earth's deep-sea sharks, starless organisms will utilize sensitive electroreceptors to detect the micro-volt electrical fields generated by the muscular movements or chemical ion currents of neighboring life forms.

8.2 Metabolic Adaptation & Energetic Constraints

Due to the rigid energy limitations imposed by a chemosynthetic baseline, complex multicellular life forms in a rogue ocean will exhibit exceptionally low metabolic rates. Growth, reproduction, and locomotion will operate on protracted, multi-decade timelines.

To minimize energy consumption, organisms will evolve extreme longevity, highly efficient closed-loop internal nutrient recycling systems, and soft, gelatinous structural forms lacking heavy, energy-expensive calcified skeletons. The ecosystem will resemble a slow-motion biosphere, optimized for near-perfect thermodynamic conservation.

9. Advanced Speculative Possibilities & Technosignatures

While the existence of microbial and basic multicellular life within rogue oceans is firmly supported by geochemical scaling laws, the emergence of technological species within these starless environments enters the domain of carefully constrained scientific speculation.

9.1 Geothermal Technological Civilizations

Could a technological civilization emerge inside a starless, subglacial world? The path to technology would be radically non-human. Deprived of an open atmosphere and fire, these species could not utilize combustion or standard metallurgy. Instead, early industrial development would rely on controlling hydrothermal flow channels and localized volcanic heat vents.

Material science would likely center on biochemical polymers, harvested ice-matrix composites, and ceramics formed via extreme hydrostatic pressures. Energy extraction would utilize highly optimized closed-loop geothermal heat exchangers, harvesting electricity directly from the stark temperature differential between the hot hydrothermal vent fields (~600 K) and the ambient ocean water (~273 K).

9.2 Traveling Worlds and Planetary Engineering

On a grand cosmic scale, highly advanced interstellar civilizations might recognize rogue planets not as barren obstacles, but as the ultimate interstellar migration vessels. A star-bound planet is perpetually exposed to severe space weather, stellar flares, and eventual stellar death. A massive rogue planet, by contrast, is completely immune to stellar cataclysms.

As proposed by recent technosignature studies, a civilization could deliberately engineer a rogue planet into a mobile traveling world (Romanovskaya, 2022). By deploying massive fusion reactors or artificial planetary magnetic fields, an advanced species could regulate the internal thermal engine, utilizing the rogue world's deep ice shell as an absolute defense against cosmic radiation during a multi-million-year trek across the interstellar void.

10. Philosophical & Cosmic Implications

The realization that habitable environments can exist completely independent of stars fundamentally alters our understanding of astrobiology and the distribution of life in the universe.

10.1 Redefining the Circumstellar Paradigm

For over a century, the search for life has focused entirely on discovering Earth twins orbiting Sun-like stars. The Steppenwolf and primordial hydrogen models break this constraint, proving that **Habitable Zone ≠ Only Region Around Stars**. If habitability is an intrinsic planetary property fueled by internal geological energy rather than external solar irradiance, then the total volume of habitable real estate in the universe increases by orders of magnitude.

10.2 Resolution of the Fermi Paradox

This paradigm shift provides a compelling, elegant potential solution to the Fermi Paradox ("Where is everyone?"). If the vast majority of biospheres—and potentially the vast majority of intelligent civilizations—originate deep inside rogue planets beneath hundreds of kilometers of impenetrable ice, their entire cosmic perspective would be completely internal.

A species evolving inside a subglacial ocean has no natural visual access to the stars, no initial concept of space, and no capacity to broadcast or receive electromagnetic radio signals through a global water-and-ice radio shield. They may build highly complex, advanced informational societies entirely within their internal abyssal ecosystems, completely invisible to our standard astronomical SETI sweeps. The universe may contain an active, ubiquitous network of life hidden away inside frozen, wandering worlds drifting silently through the dark interstellar void.

References

Abbot, D. S., & Switzer, E. R. (2011). THE STEPPENWOLF: A PROPOSAL FOR A HABITABLE PLANET IN INTERSTELLAR SPACE. *The Astrophysical Journal*, 735(2), L27. https://doi.org/10.1088/2041-8205/735/2/l27
Cited by: 78

Balbi, A., Hami, M., & Kovačević, A. B. (2020). The Habitability of the Galactic Bulge. *arXiv e-prints*, arXiv:1008.0132. https://doi.org/10.3390/life10080132
Cited by: 28

Romanovskaya, I. K. (2022). Migrating extraterrestrial civilizations and interstellar colonization: implications for SETI and SETA. *International Journal of Astrobiology*, 21(3), 163–187. https://doi.org/10.1017/s1473550422000143
Cited by: 14

Sadlok, G. (2020). On A Hypothetical Mechanism of Interstellar Life Transfer Through Nomadic Objects. *Origins of Life and Evolution of Biospheres*, 50(2), 87–96. https://doi.org/10.1007/s11084-020-09591-z
Cited by: 9

Schulze-Makuch, D., & Fairén, A. G. (2021). Evaluating the Microbial Habitability of Rogue Planets and Proposing Speculative Scenarios on How They Might Act as Vectors for Panspermia. *Life*, 11(8), 833. https://doi.org/10.3390/life11080833
Cited by: 8

Strigari, L. E., Barnabe, M., Marshall, P. J., & Blandford, R. D. (2012). Nomads of the Galaxy. *Monthly Notices of the Royal Astronomical Society*, 423(2), 1856–1865. https://doi.org/10.1111/j.1365-2966.2012.21009.x
Cited by: 86

Zhang, X. (2020). Atmospheric regimes and trends on exoplanets and brown dwarfs. *Research in Astronomy and Astrophysics*, 20(7), 099. https://doi.org/10.1088/1674-4527/20/7/99
Cited by: 109